

# Parents - Are Your Kids Ready for School?

## Required Immunizations for School Year 2013-2014



Parent/Guardian Resource

	<b>Hepatitis B</b>	<b>DTaP/Td/Tdap*</b> (Diphtheria, Tetanus, Pertussis)	<b>Polio*</b>	<b>MMR</b> (Measles, Mumps, Rubella)	<b>Varicella</b> (Chickenpox)
<b>Kindergarten – 5<sup>th</sup> Grade</b>	3 doses	5 doses	4 doses	2 doses	2 doses OR Doctor verifies disease
<b>6<sup>th</sup> Grade</b>	3 doses	5 doses DTaP  <i>AND</i>  1 dose Tdap	4 doses	2 doses	1 dose OR Parent verifies disease
<b>7<sup>th</sup> – 12<sup>th</sup> Grade</b>	3 doses	5 doses DTaP  <i>AND</i>  1 dose Tdap	4 doses	2 doses	Recommended, but not required.

\*Vaccine doses required may be fewer than listed.

- Students must meet minimum intervals and ages to be in compliance with the requirements. Talk to your healthcare provider or school staff if you have questions about school immunization requirements.
- Find information on other recommended vaccines not required for school: <http://www.immunize.org/cdc/schedules/>

# Padres de familia

## ¿Estan sus hijos listos para la escuela?



Vacunas requeridas para el ciclo escolar 2013-2014

Recursos para padres de familia y tutores legales

	<b>Hepatitis B</b>	<b>DTaP/Td/Tdap*</b> (Difteria, Tétanos, Tos Ferina)	<b>Polio*</b>	<b>MMR</b> (SPR-Sarampión, Paperas, Rubeola)	<b>Varicela</b>
<b>Kínder – 5<sup>o</sup> Grado</b>	3 dosis	5 dosis	4 dosis	2 dosis	2 dosis O El doctor verifica la enfermedad
<b>6<sup>o</sup> Grado</b>	3 dosis	5 dosis DTaP  Y  1 dosis de Tdap	4 dosis	2 dosis	1 dosis O Padres verifican la enfermedad
<b>7<sup>o</sup> – 12<sup>o</sup> Grado</b>	3 dosis	5 dosis DTaP  Y  1 dosis de Tdap	4 dosis	2 dosis	Recomendada pero no requerida.

\* Las dosis de vacunas requeridas podrían ser menos de las que aquí se enlistan.

- Los estudiantes deben cumplir con los intervalos mínimos y edades para estar al día con los requisitos de vacunación. Si usted tiene preguntas sobre los requisitos de vacunación para el ingreso escolar, hable con su proveedor de salud o personal escolar capacitado.
- Encuentre información sobre otras vacunas recomendadas pero no requeridas para el ingreso escolar en: [www.immunize.org/cdc/schedules/](http://www.immunize.org/cdc/schedules/)



## VACCINES REQUIRED FOR SCHOOL ATTENDANCE, GRADES K-12

July 1, 2013 – June 30, 2014

VACCINE	Kindergarten-2 <sup>nd</sup> Grade	3 <sup>rd</sup> -5 <sup>th</sup> Grade	6 <sup>th</sup> Grade	7 <sup>th</sup> -12 <sup>th</sup> Grade
<b>Hepatitis B</b>	<b>3 doses</b> See minimum intervals on page 2 Dose 3 must be given $\geq 24$ weeks of age			<b>3 doses</b> See minimum intervals on page 2 Dose 3 must be given $\geq 4$ months of age
<b>Diphtheria, Tetanus, and Pertussis (DTaP/DT/Td/Tdap)</b>	<b>5 doses</b> (4 doses required <b>IF</b> 4 <sup>th</sup> dose given $\geq 4^{\text{th}}$ birthday)  <i>plus</i>  <b>1 dose Tdap required for 6<sup>th</sup>-12<sup>th</sup> grade IF <math>\geq 11</math> years old</b>			
<b>Polio (IPV or OPV)</b>	<b>4 doses</b> (3 doses only <b>IF</b> 3 <sup>rd</sup> dose given $\geq 4^{\text{th}}$ birthday) <ul style="list-style-type: none"> <li>The final dose given on or after August 7, 2009 must be given at a minimum of 4 years of age AND a minimum interval of 6 months from the previous dose.</li> </ul>	<b>4 doses</b> (3 doses only <b>IF</b> 3 <sup>rd</sup> dose given $\geq 4^{\text{th}}$ birthday)		
<b>Measles, Mumps, and Rubella</b>	<b>2 doses</b>			
<b>Varicella</b>	<b>2 doses</b>  <b>OR</b>  Healthcare provider verifies disease		<b>1 dose</b>  <b>OR</b>  Parent verifies disease	<b>Recommended</b> , but not required.

- Look at the Minimum Age and Interval Table on page 2 for recommended minimum age and spacing information.
- Review the Individual Vaccine Requirements Summary for more detailed information: [www.doh.wa.gov/Portals/1/Documents/Pubs/348-284-IndividualVaccineRequirements.pdf](http://www.doh.wa.gov/Portals/1/Documents/Pubs/348-284-IndividualVaccineRequirements.pdf)

## Minimum Age & Interval for Valid Vaccine Doses

Vaccine	Dose #	Minimum Age	Minimum Interval Between Doses	Notes
Hepatitis B <b>HepB</b>	Dose 1	Birth	4 weeks between Dose 1 & 2 (K-12 <sup>th</sup> )	<ul style="list-style-type: none"> <li>▪ Note minimum age and interval changes for 2013-14 school year.</li> <li>▪ 2 doses valid if adult Recombivax HB<sup>®</sup> given between ages 11 and 15 and doses separated by <math>\geq 4</math> months.</li> </ul>
	Dose 2	4 weeks	8 weeks between Dose 2 & 3 (K-12 <sup>th</sup> )	
	Dose 3	24 weeks (K-6 <sup>th</sup> ) 4 months (7 <sup>th</sup> -12 <sup>th</sup> )	16 weeks between Dose 1 & 3 (K-6 <sup>th</sup> ) No minimum interval between Dose 1 & 3 (7 <sup>th</sup> -12 <sup>th</sup> )	
Diphtheria, Tetanus, and Pertussis <b>DTaP/DT</b>	Dose 1	6 weeks	4 weeks between Dose 1 & 2	<ul style="list-style-type: none"> <li>▪ DTaP: for children through age 6.</li> <li>▪ Recommended to have 6 months between Dose 3 and Dose 4, but <math>\geq 4</math> months acceptable.</li> </ul>
	Dose 2	10 weeks	4 weeks between Dose 2 & 3	
	Dose 3	14 weeks	6 months between Dose 3 & 4	
	Dose 4	12 months	6 months between Dose 4 & 5	
	Dose 5	4 years	–	
Tetanus, Diphtheria, and Pertussis <b>Tdap</b>	Dose 1	10 years (minimum age depends on vaccine brand)	–	<ul style="list-style-type: none"> <li>▪ Boostrix<sup>®</sup>: licensed for <math>\geq 10</math> year olds; Adacel<sup>®</sup>: licensed for <math>\geq 11</math> year olds.</li> <li>▪ Can be given regardless of the interval between DTaP or Td.</li> <li>▪ Students 7-10 years of age not fully immunized with DTaP or Td should get one Tdap followed by additional doses of Td if needed.</li> </ul>
Tetanus and Diphtheria <b>Td</b>	Dose 1	7 years	5 years	<ul style="list-style-type: none"> <li>▪ Td: for children <math>\geq 7</math> years of age.</li> <li>▪ 3 doses of Td required, if starting series <math>\geq 7</math> years, with a single dose of Tdap preferred as the first dose.</li> </ul>
Polio <b>IPV or OPV</b>	Dose 1	6 weeks	4 weeks between Dose 1 & 2	<ul style="list-style-type: none"> <li>▪ Not required for students 18 years and older.</li> <li>▪ If a student got all doses before August 7, 2009: 4 week minimum interval must separate all doses and minimum age must be <math>\geq 18</math> weeks.</li> </ul>
	Dose 2	10 weeks	4 weeks between Dose 2 & 3	
	Dose 3	14 weeks	6 months between Dose 3 & 4	
	Dose 4	4 years	–	
Measles, Mumps, and Rubella <b>MMR</b>	Dose 1	12 months	4 weeks between Dose 1 & 2	<ul style="list-style-type: none"> <li>▪ MMRV (MMR + varicella) may be used instead of separate MMR and varicella vaccines.</li> <li>▪ 4-day grace <b>DOES</b> apply between doses of the same live vaccine such as MMR/MMR or MMRV/MMRV. The 4 day grace period <b>DOES NOT</b> apply between Dose 1 and Dose 2 of different live vaccines, such as between MMR and Varicella or between MMR and live flu vaccine.</li> </ul>
	Dose 2	13 months	–	
Varicella (chickenpox) <b>VAR</b>	Dose 1	12 months	3 months between Dose 1 & 2	<ul style="list-style-type: none"> <li>▪ Recommended: 3 months between varicella doses, but <math>\geq 28</math> days acceptable.</li> <li>▪ Must get the same day as MMR <b>OR</b> <math>\geq 28</math> days apart. (4-day grace <b>DOES NOT</b> apply).</li> <li>▪ 4-day grace <b>DOES</b> apply between doses of the same live vaccine such as VAR and VAR).</li> </ul>
	Dose 2	15 months	–	

If you have a disability and need this document in another format, please call 1-800-525-0127 (TDD/TTY call 711)

DOH 348-051 January 2013